

## **HPTLC-ESI-MS/MS for identifying neutral lipids, sphingolipids and phospholipids in complex samples**

**CEBOLLA V**

vcebolla@icb.csic.es

Cebolla V<sup>1</sup>, Lapieza M<sup>1</sup>, Jarne C<sup>1</sup>, Saviron M<sup>2</sup>, Orduna J<sup>3</sup>, Membrado L<sup>1</sup>, Morlock GE<sup>5</sup>,  
Jungas C<sup>6</sup>, Garriga R<sup>4</sup>, Vela J<sup>4</sup>

<sup>1</sup>CSIC, <sup>2</sup>CEQMA-CSIC, <sup>3</sup>CSIC-ICMA and <sup>4</sup>University of Zaragoza, Zaragoza, Spain,

<sup>4</sup>Justus Liebig University Giessen, Chair of Food Science, Giessen, Germany,

<sup>5</sup>CEA-CNRS-Aix Marseille University, Marseille, France

HPTLC is well-adapted to providing lipid-class separations, as an assistant technique of HPLC. From this perspective, it can complement traditional LC-mass spectrometric approaches in a unique way. As the zones of interest on the plate can selectively be transferred to the ESI-MS instrument via an elution head-based interface, a rapid, relevant structural information about molecular species within each lipid classes from complex samples can be obtained.

Separation using AMD2 provides lipid-classes as narrow peaks enough to ensure a direct elution and transfer of the plate zones to obtain both composition profiles of each class by ESI-MS, and identification of individual lipids and molecular species belonging to each separated lipid class by MS/MS (MS<sup>n</sup>) and HRMS. The respective sodium adducts of the above-mentioned lipid classes were fragmented in the positive ion mode using an ion-trap technology. The sodium remained the charge of their fragment ions, thus being useful for their structural identification by MS/MS (MS<sup>n</sup>) through further fragmentation.

This work exemplarily focuses on profiling and identification of neutral lipids (NLs), sphingolipids (SLs) and phospholipids (PLs) in three analytical cases:

- (1) mono- (MAGs) and diacylglycerides (DAGs) in positive ion mode and fatty acids (FAs) in negative ion mode as impurities in a fatty acid methyl ester-based biodiesel
- (2) molecular species of neutral sphingolipids (SL), such as sphingomyelins (SMs) and globotriaosylceramides (Gb3), in human plasma
- (3) phosphatidylcholines (PCs), phosphatidylethanolamines (PEs), phosphatidylglycerols (PGs) and cardiolipines (CLs) associated to membrane proteins of photosynthetic purple bacteria.